arXiv:1606.03791v2 [math.NT] 2 Apr 2017

UNIVOQUE BASES AND HAUSDORFF DIMENSIONS

DERONG KONG, WENXIA LI, FAN LÜ, AND MARTIJN DE VRIES

Abstract. Given a positive integer M and a real number $q > 1$, a *q-expansion* of a real number x is a sequence $(c_i) = c_1c_2\cdots$ with $(c_i) \in \{0,\dots,M\}^\infty$ such that

$$x = \sum_{i=1}^{\infty} c_i q^{-i}.$$

It is well known that if $q \in (1, M+1]$, then each $x \in I_q := [0, M/(q-1)]$ has a q-expansion. Let $\mathcal{U} = \mathcal{U}(M)$ be the set of *univoque bases* $q > 1$ for which 1 has a unique q-expansion. The main object of this paper is to provide new characterizations of $\mathcal{U}$ and to show that the Hausdorff dimension of the set of numbers $x \in I_q$ with a unique q-expansion changes the most if q "crosses" a univoque base.

Denote by $\mathcal{B}_2 = \mathcal{B}_2(M)$ the set of $q \in (1, M+1]$ such that there exist numbers having precisely two distinct q-expansions. As a by-product of our results, we obtain an answer to a question of Sidorov (2009) and prove that

$$\dim_H(\mathcal{B}_2 \cap (q', q'+\delta)) > 0 \quad \text{for any} \quad \delta > 0,$$

where $q' = q'(M)$ is the Komornik-Loreti constant.

1. Introduction

Non-integer base expansions have received much attention since the pioneering works of Rényi [26] and Parry [25]. Given a positive integer M and a real number $q \in (1, M+1]$, a sequence $(d_i) = d_1d_2\cdots$ with *digits* $d_i \in \{0, 1, \dots, M\}$ is called a *q-expansion of x* or an *expansion of x in base q* if

$$x = \pi_q((d_i)) := \sum_{i=1}^{\infty} \frac{d_i}{q^i}.$$

It is well known that each $x \in I_q := [0, M/(q-1)]$ has a q-expansion. One such expansion - the *greedy q-expansion* - can be obtained by performing the so called *greedy algorithm* of Rényi which is defined recursively as follows: if $d_1, \dots, d_{n-1}$ is already defined (no condition if $n = 1$), then d_n is the largest element of $\{0, \dots, M\}$ satisfying $\sum_{i=1}^{n} d_i q^{-i} \le x$.

Date: April 4, 2017.

1991 *Mathematics Subject Classification.* 11A63 and 37B10 and 28A78.

Key words and phrases. univoque bases and univoque sets and Hausdorff dimensions and generalized Thue-Morse sequences.

Equivalently, (d_i) is the greedy q-expansion of $\sum_{i=1}^{\infty} d_i q^{-i}$ if and only if $\sum_{i=n+1}^{\infty} d_i q^{-i+n} < 1$ whenever $d_n < M$, $n = 1, 2, \ldots$. Hence if $1 < q < r \leq M+1$, then the greedy q-expansion of a number $x \in I_q$ is also the greedy expansion in base r of a number in I_r.

Let $\mathcal{U}_q$ be the *univoque set* consisting of numbers $x \in I_q$ such that x has a unique q-expansion, and let $\mathcal{U}_q'$ be the set of corresponding expansions. Note that a sequence (c_i) belongs to $\mathcal{U}_q'$ if and only if both the sequences (c_i) and $(M - c_i) := (M - c_1)(M - c_2) \cdots$ are greedy q-expansions, hence $\mathcal{U}_q' \subseteq \mathcal{U}_r'$ whenever $1 < q < r \leq M+1$. Many works are devoted to the univoque sets $\mathcal{U}_q$ (see, e.g., [11, 13, 15]). Recently, de Vries and Komornik investigated their topological properties in [9]. Komornik et al. considered their Hausdorff dimension in [17], and showed that the dimension function $D : q \mapsto \dim_H \mathcal{U}_q$ behaves like a Devil's staircase on $(1, M+1]$. For more information on the univoque set $\mathcal{U}_q$ we refer to the survey paper [16] and the references therein.

There is an intimate connection between the set $\mathcal{U}_q$ and the set of *univoque bases* $\mathcal{U} = \mathcal{U}(M)$ consisting of numbers $q > 1$ such that 1 has a unique q-expansion over the alphabet $\{0, 1, \ldots, M\}$. For instance, it was shown in [9] that $\mathcal{U}_q$ is closed if and only if q does not belong to the set $\overline{\mathcal{U}}$. It is well-known that $\mathcal{U}$ is a Lebesgue null set of full Hausdorff dimension (cf. [7, 12, 17]). Moreover, the smallest element of $\mathcal{U}$ is the *Komornik-Loreti constant* (cf. [18, 19])

$$q' = q'(M),$$

while the largest element of $\mathcal{U}$ is (of course) $M+1$. Recently, Komornik and Loreti showed in [20] that its closure $\overline{\mathcal{U}}$ is a *Cantor set* (see also, [10]), i.e., a nonempty closed set having neither isolated nor interior points. Writing the open set $(1, M+1] \setminus \overline{\mathcal{U}} = (1, M+1) \setminus \overline{\mathcal{U}}$ as the disjoint union of its connected components, i.e.,

$$(1, M+1] \setminus \overline{\mathcal{U}} = (1, q') \cup \bigcup (q_0, q_0^*), \tag{1.1}$$

the left endpoints q_0 in (1.1) run over the whole set $\overline{\mathcal{U}} \setminus \mathcal{U}$, and the right endpoints q_0^* run through a subset of $\mathcal{U}$ (cf. [9]). Furthermore, each left endpoint q_0 is algebraic, while each right endpoint $q_0^* \in \mathcal{U}$ is transcendental (cf. [21]).

De Vries showed in [8], roughly speaking, that the sets $\mathcal{U}_q'$ change the most if we cross a univoque base. More precisely, it was shown that $q \in \mathcal{U}$ if and only if $\mathcal{U}_r' \setminus \mathcal{U}_q'$ is uncountable for each $r \in (q, M+1]$ and $r \in \overline{\mathcal{U}}$ if and only if $\mathcal{U}_r' \setminus \mathcal{U}_q'$ is uncountable for each $q \in (1, r)$.

The main object of this paper is to provide similar characterizations of $\mathcal{U}$ and $\overline{\mathcal{U}}$ in terms of the Hausdorff dimension of the sets $\mathcal{U}_r' \setminus \mathcal{U}_q'$ after a natural projection. Furthermore, we characterize the sets $\mathcal{U}$ and $\overline{\mathcal{U}}$ by looking at the Hausdorff dimensions of $\mathcal{U}$ and $\overline{\mathcal{U}}$ locally.

Theorem 1.1. *Let $q \in (1, M+1]$. The following statements are equivalent.*

(i) $q \in \mathcal{U}$.
(ii) $\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) > 0$ *for any* $r \in (q, M+1]$.
(iii) $\dim_H \mathcal{U} \cap (q, r) > 0$ *for any* $r \in (q, M+1]$.

Theorem 1.2. *Let* $q \in (1, M+1]$. *The following statements are equivalent.*

(i) $q \in \overline{\mathcal{U}} \setminus (\bigcup \{q_0^*\} \cup \{q'\})$.
(ii) $\dim_H \pi_{M+1}(\mathcal{U}'_q \setminus \mathcal{U}'_p) > 0$ *for any* $p \in (1, q)$.
(iii) $\dim_H \mathcal{U} \cap (p, q) > 0$ *for any* $p \in (1, q)$.

It follows at once from Theorems 1.1 and 1.2 that $\mathcal{U}$ (or, equivalently, $\overline{\mathcal{U}}$) does not contain isolated points.

We remark that the projection map π_{M+1} in Theorem 1.1 (ii) can be replaced by π_ρ for any $r \le \rho \le M+1$. Similarly, the projection map π_{M+1} in Theorem 1.2 (ii) can also be replaced by π_ρ with $q \le \rho \le M+1$. We also point out that Theorems 1.1 and 1.2 strengthen the main result of [8] where the cardinality of the sets $\mathcal{U}'_q \setminus \mathcal{U}'_p$ with $1 < p < q \le M+1$ was determined.

Let $\mathcal{B}_2$ be the set of bases $q \in (1, M+1]$ for which there exists a number $x \in [0, M/(q-1)]$ having exactly two q-expansions. It was asked by Sidorov [27] whether $\dim_H \mathcal{B}_2 \cap (q', q' + \delta) > 0$ for any $\delta > 0$, where q' is the Komornik-Loreti constant. Since $\mathcal{U} \subseteq \mathcal{B}_2$ (see [27, Lemma 3.1][1]), Theorem 1.1 answers this question in the affirmative.

Corollary 1.3. $\dim_H \mathcal{B}_2 \cap (q', q' + \delta) > 0$ *for any* $\delta > 0$.

The rest of the paper is arranged as follows. In Section 2 we recall some properties of unique q-expansions. The proof of Theorems 1.1 and 1.2 will be given in Section 3.

2. Preliminaries

In this section we recall some properties of the univoque set $\mathcal{U}_q$. Throughout this paper, a *sequence* $(d_i) = d_1 d_2 \cdots$ is an element of $\{0, \dots, M\}^\infty$ with each digit d_i belonging to the *alphabet* $\{0, \dots, M\}$. Moreover, for a *word* $\mathbf{c} = c_1 \cdots c_n$ we mean a finite string of digits with each digit c_i from $\{0, \dots, M\}$. For two words $\mathbf{c} = c_1 \cdots c_n$ and $\mathbf{d} = d_1 \cdots d_m$ we denote by $\mathbf{cd} = c_1 \cdots c_n d_1 \cdots d_m$ the concatenation of the two words. For an integer $k \ge 1$ we denote by $\mathbf{c}^k$ the k-times concatenation of $\mathbf{c}$ with itself, and by $\mathbf{c}^\infty$ the infinite repetition of $\mathbf{c}$.

For a sequence (d_i) we denote its *reflection* by $\overline{(d_i)} := (M-d_1)(M-d_2)\cdots$. Accordingly, for a word $\mathbf{c} = c_1 \cdots c_n$ we denote its reflection by $\overline{\mathbf{c}} := (M-c_1)\cdots(M-c_n)$. If $c_n < M$ we denote by $\mathbf{c}^+ := c_1 \cdots c_{n-1}(c_n+1)$. If $c_n > 0$ we write $\mathbf{c}^- := c_1 \cdots c_{n-1}(c_n-1)$.

[1]This also follows directly from the observation that q^{-1} has exactly two q-expansions whenever $q \in \mathcal{U}$.

We will use systematically the lexicographic ordering $<, \leq, >$ and $\geq$ between sequences and between words. For two sequences $(c_i), (d_i) \in \{0, 1, \ldots, M\}^{\infty}$ we say that $(c_i) < (d_i)$ if there exists an integer $n \geq 1$ such that $c_1 \ldots c_{n-1} = d_1 \ldots d_{n-1}$ and $c_n < d_n$. Furthermore, we write $(c_i) \leq (d_i)$ if $(c_i) < (d_i)$ or $(c_i) = (d_i)$. Similarly, we say $(c_i) > (d_i)$ if $(d_i) < (c_i)$, and $(c_i) \geq (d_i)$ if $(d_i) \leq (c_i)$. We extend this definition to words in the obvious way. For example, for two words $\mathbf{c}$ and $\mathbf{d}$ we write $\mathbf{c} < \mathbf{d}$ if $\mathbf{c}0^{\infty} < \mathbf{d}0^{\infty}$.

A sequence is called *finite* if it has a last nonzero element. Otherwise it is called *infinite*. So $0^{\infty} := 00\cdots$ is considered to be infinite. For $q \in (1, M+1]$ we denote by

$$\alpha(q) = (\alpha_i(q))$$

the *quasi-greedy* q-expansion of 1 (cf. [6]), i.e., the lexicographically largest *infinite* q-expansion of 1. Let $\beta(q) = (\beta_i(q))$ be the *greedy* q-expansion of 1 (cf. [25]), i.e., the lexicographically largest q-expansion of 1. For convenience, we set $\alpha(1) = 0^{\infty}$ and $\beta(1) = 10^{\infty}$, even though $\alpha(1)$ is not a 1-expansion of 1.

Moreover, we endow the set $\{0, \ldots, M\}$ with the discrete topology and the set of all possible sequences $\{0, 1, \ldots, M\}^{\infty}$ with the Tychonoff product topology.

The following properties of $\alpha(q)$ and $\beta(q)$ were established in [25], see also [4].

Lemma 2.1.

(i) *The map $q \mapsto \alpha(q)$ is an increasing bijection from $[1, M+1]$ onto the set of all* infinite *sequences (α_i) satisfying*

$$\alpha_{n+1}\alpha_{n+2}\cdots \leq \alpha_1\alpha_2\cdots \quad \text{whenever} \quad \alpha_n < M.$$

(ii) *The map $q \mapsto \beta(q)$ is an increasing bijection from $[1, M+1]$ onto the set of all sequences (β_i) satisfying*

$$\beta_{n+1}\beta_{n+2}\cdots < \beta_1\beta_2\cdots \quad \text{whenever} \quad \beta_n < M.$$

Lemma 2.2.

(i) *$\beta(q)$ is infinite if and only if $\beta(q) = \alpha(q)$.*
(ii) *If $\beta(q) = \beta_1\cdots\beta_m 0^{\infty}$ with $\beta_m > 0$, then $\alpha(q) = (\beta_1\cdots\beta_m^-)^{\infty}$.*
(iii) *The map $q \mapsto \alpha(q)$ is left-continuous, while the map $q \mapsto \beta(q)$ is right-continuous.*

In order to investigate the unique expansions we need the following lexicographic characterization of $\mathcal{U}_q'$ (cf. [4]).

Lemma 2.3. *Let $q \in (1, M+1]$. Then $(d_i) \in \mathcal{U}_q'$ if and only if*

$$\begin{cases} d_{n+1}d_{n+2}\cdots < \overline{\alpha_1(q)\alpha_2(q)\cdots} & \text{whenever} \quad d_n < M, \\ d_{n+1}d_{n+2}\cdots > \overline{\alpha_1(q)\alpha_2(q)\cdots} & \text{whenever} \quad d_n > 0. \end{cases}$$

Note that $q \in \mathcal{U}$ if and only if $\alpha(q)$ is the unique q-expansion of 1. Then Lemma 2.3 yields a characterization of $\mathcal{U}$ (see also, [11] and [19]).

Lemma 2.4. *Let $q \in (1, M+1)$. Then $q \in \mathcal{U}$ if and only if $\alpha(q) = (\alpha_i(q))$ satisfies*

$$\overline{\alpha(q)} < \alpha_{n+1}(q)\alpha_{n+2}(q)\cdots < \alpha(q) \quad \text{for all} \quad n \geq 1.$$

Consider a connected component (q_0, q_0^*) of $(q', M+1)\setminus \overline{\mathcal{U}}$ as in (1.1). Then there exists a (unique) word $\mathbf{t} = t_1 \cdots t_p$ such that (cf. [9, 21])

$$\alpha(q_0) = \mathbf{t}^\infty \quad \text{and} \quad \alpha(q_0^*) = \lim_{n\to\infty} g^n(\mathbf{t}),$$

where $g^n = \underbrace{g \circ \cdots \circ g}_{n}$ denotes the n-fold composition of g with itself, and

$$g(\mathbf{c}) := \mathbf{c}^+\overline{\mathbf{c}^+} \quad \text{for any word} \quad \mathbf{c} = c_1 \ldots c_k \text{ with } c_k < M. \tag{2.1}$$

We point out that the word $\mathbf{t} = t_1 \ldots t_p$ in the definitions of $\alpha(q_0)$ and $\alpha(q_0^*)$ is called an *admissible block* in [21, Definition 2.1] which satisfies the following lexicographical inequalities: $t_p < M$ and for any $1 \leq i \leq p$ we have

$$\overline{t_1 \cdots t_p} \leq t_i \cdots t_p t_1 \cdots t_{i-1} \quad \text{and} \quad t_i \cdots t_p\, \overline{t_1 \ldots t_{i-1}} \leq t_1 \ldots t_p^+.$$

We also mention that the limit $\lim_{n\to\infty} g^n(\mathbf{t})$ stands for the infinite sequence beginning with $\mathbf{t}^+\overline{\mathbf{t}}\,\overline{\mathbf{t}^+}\mathbf{t}^+\,\overline{\mathbf{t}^+}\mathbf{t}\,\mathbf{t}^+\overline{\mathbf{t}}\cdots$, and the existence of this limit was shown by Allouche [1].

In this case (q_0, q_0^*) is called the *connected component generated by* $\mathbf{t}$. The closed interval $[q_0, q_0^*]$ is the so called *admissible interval generated by* $\mathbf{t}$ (see [21, Definition 2.4]). Furthermore, the sequence

$$\alpha(q_0^*) = \lim_{n\to\infty} g^n(\mathbf{t}) = \mathbf{t}^+\,\overline{\mathbf{t}}\,\overline{\mathbf{t}^+}\,\mathbf{t}^+\,\overline{\mathbf{t}^+}\,\mathbf{t}\,\mathbf{t}^+\overline{\mathbf{t}}\cdots$$

is a generalized Thue-Morse sequence (cf. [21, Definition 2.2], see also [3]).

The following lemma for the generalized Thue-Morse sequence $\alpha(q_0^*)$ was established in [21, Lemma 4.2].

Lemma 2.5. *Let $(q_0, q_0^*) \subset (q', M+1)\setminus\overline{\mathcal{U}}$ be a connected component generated by $t_1 \cdots t_p$. Then the sequence $(\theta_i) = \alpha(q_0^*)$ satisfies*

$$\overline{\theta_1 \cdots \theta_{2^n p - i}} < \theta_{i+1} \cdots \theta_{2^n p} \leq \theta_1 \cdots \theta_{2^n p - i}$$

for any $n \geq 0$ and any $0 \leq i < 2^n p$.

Finally, we recall some topological properties of $\mathcal{U}$ and $\overline{\mathcal{U}}$ which were essentially established in [9, 20] (see also, [10]).

Lemma 2.6.

(i) *If $q \in \mathcal{U}$, then there exists a* decreasing *sequence (r_n) of elements in $\bigcup\{q_0^*\}$ that converges to q as $n \to \infty$;*

(ii) *If $q \in \overline{\mathcal{U}} \setminus (\bigcup\{q_0^*\} \cup \{q'\})$, then there exists an* increasing *sequence (p_n) of elements in $\bigcup\{q_0^*\}$ that converges to q as $n \to \infty$.*

We remark here that the bases q_0^* are called *de Vries-Komornik numbers* which were shown to be transcendental in [21]. By Lemma 2.6 it follows that the set of de Vries-Komornik numbers is dense in $\overline{\mathcal{U}}$.

3. Proofs of Theorems 1.1 and 1.2

3.1. **Proof of Theorem 1.1 for (i) $\Leftrightarrow$ (ii).** For each connected component (q_0, q_0^*) of $(q', M+1) \setminus \overline{\mathcal{U}}$ we construct a sequence of bases (r_n) in $\mathcal{U}$ strictly decreasing to q_0^*.

Lemma 3.1. *Let $(q_0, q_0^*) \subset (q', M+1) \setminus \overline{\mathcal{U}}$ be a connected component generated by $t_1 \cdots t_p$, and let $(\theta_i) = \alpha(q_0^*)$. Then for each $n \geq 1$, the number $r_n \in \mathcal{U}$ determined by*

$$\alpha(r_n) = \beta(r_n) = \theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty,$$

belongs to $\mathcal{U}$. Furthermore, (r_n) is a strictly decreasing sequence that converges to q_0^*.

Proof. Using (2.1) one may verify that the sequence (θ_i) satisfies

$$\theta_{2^n p+k} = \overline{\theta_k} \quad \text{for all} \quad 1 \leq k < 2^n p; \quad \theta_{2^{n+1}p} = \overline{\theta_{2^n p}}^+$$

for all $n \geq 0$. Now fix $n \geq 1$. We claim that

$$\sigma^i\left(\theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty\right) < \theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty \tag{3.1}$$

for all $i \geq 1$, where σ is the left shift on $\{0, \ldots, M\}^\infty$ defined by $\sigma((c_i)) = (c_{i+1})$. By periodicity it suffices to prove (3.1) for $0 < i < 2^{n+1}p$. We distinguish between the following three cases: (I) $0 < i < 2^n p$; (II) $i = 2^n p$; (III) $2^n p < i < 2^{n+1}p$.

Case (I). $0 < i < 2^n p$. Then by Lemma 2.5 it follows that

$$\theta_{i+1} \cdots \theta_{2^n p} \leq \theta_1 \cdots \theta_{2^n p - i}$$

and

$$\theta_{2^n p+1} \cdots \theta_{2^n p+i} = \overline{\theta_1 \cdots \theta_i} < \theta_{2^n p-i+1} \cdots \theta_{2^n p}.$$

This implies (3.1) for $0 < i < 2^n p$.

Case (II). $i = 2^n p$. Note by [19] that $\alpha_1(q') = [M/2] + 1$ (see also, [5]), where $[y]$ denotes the integer part of a real number y. Then by using $q_0^* > q'$ in Lemma 2.1 we have

$$\theta_1 = \alpha_1(q_0^*) \geq \alpha_1(q') > \overline{\alpha_1(q')} \geq \overline{\theta_1}.$$

This, together with $n \geq 1$, implies

$$\theta_{2^n p+1} \cdots \theta_{2^{n+1}p} = \overline{\theta_1 \cdots \theta_{2^n p}}^{+} < \theta_1 \cdots \theta_{2^n p}.$$

So, (3.1) holds true for $i = 2^n p$.

Case (III). $2^n p < i < 2^{n+1}p$. Write $j = i - 2^n p$. Then $0 < j < 2^n p$. Once again, we infer from Lemma 2.5 that

$$\theta_{i+1} \cdots \theta_{2^{n+1}p} = \overline{\theta_{j+1} \cdots \theta_{2^n p}}^{+} \leq \theta_1 \cdots \theta_{2^n p - j}$$

and

$$\theta_{2^n p+1} \cdots \theta_{2^n p+j} = \overline{\theta_1 \cdots \theta_j} < \theta_{2^n p-j+1} \cdots \theta_{2^n p}.$$

This yields (3.1) for $2^n p < i < 2^{n+1}p$.

Note by Lemma 2.5 that

$$\sigma^i \left(\theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty\right) > \overline{\theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty}$$

for any $i \geq 0$. Then by (3.1) and Lemma 2.4 it follows that there exists $r_n \in \mathcal{U}$ such that

$$\alpha(r_n) = \beta(r_n) = \theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty.$$

In the following we prove $r_n \searrow q_0^*$ as $n \to \infty$. For $n \geq 1$ we observe that

$$\begin{aligned}\beta(r_{n+1}) = \theta_1 \cdots \theta_{2^{n+1}p}(\theta_{2^{n+1}p+1} \cdots \theta_{2^{n+2}p})^\infty &= \theta_1 \cdots \theta_{2^n p}\overline{\theta_1 \cdots \theta_{2^n p}}^{+}\overline{\theta_1 \cdots \theta_{2^n p}} \cdots \\ &< \theta_1 \cdots \theta_{2^n p}(\overline{\theta_1 \cdots \theta_{2^n p}}^{+})^\infty = \beta(r_n).\end{aligned}$$

Then by Lemma 2.1 (ii) we have $r_{n+1} < r_n$. Note that $\beta(q_0^*) = \alpha(q_0^*) = (\theta_i)$, and

$$\beta(r_n) \to (\theta_i) = \beta(q_0^*) \quad \text{as} \quad n \to \infty.$$

Hence, we conclude from Lemma 2.2 (iii) that $r_n \searrow q_0^*$ as $n \to \infty$. □

Lemma 3.2. *Let $(q_0, q_0^*) \subset (q', M+1) \setminus \overline{\mathcal{U}}$ be a connected component generated by $t_1 \cdots t_p$, and let $(\theta_i) = \alpha(q_0^*)$. Then for any $n \geq 1$ and any $0 \leq i < 2^n p$ we have*

$$\begin{aligned}&\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\xi_n\overline{\xi_n}) < \theta_1 \cdots \theta_{2^{n+1}p-i},\\ &\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\xi_n\overline{\xi_n^-}) \leq \theta_1 \cdots \theta_{2^{n+1}p-i},\\ &\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\xi_n^-\xi_n) < \theta_1 \cdots \theta_{2^{n+1}p-i},\end{aligned} \tag{3.2}$$

and thus (by symmetry),

$$\begin{aligned}&\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\overline{\xi_n}\xi_n) < \theta_1 \cdots \theta_{2^{n+1}p-i},\\ &\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} \leq \sigma^i(\overline{\xi_n}\xi_n^-) < \theta_1 \cdots \theta_{2^{n+1}p-i},\\ &\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\overline{\xi_n^-}\,\overline{\xi_n}) < \theta_1 \cdots \theta_{2^{n+1}p-i},\end{aligned}$$

where $\xi_n := \theta_1 \cdots \theta_{2^n p}$.

Proof. By symmetry it suffices to prove (3.2).

Note that $\xi_n\overline{\xi_n} = \theta_1 \cdots \theta^-_{2^{n+1}p}$ and $\xi_n\overline{\xi_n^-} = \theta_1 \cdots \theta_{2^{n+1}p}$. Then by Lemma 2.5 it follows that

$$\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\xi_n\overline{\xi_n}) < \theta_1 \cdots \theta_{2^{n+1}p-i}$$

and

$$\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\xi_n\overline{\xi_n^-}) \le \theta_1 \cdots \theta_{2^{n+1}p-i}$$

for any $0 \le i < 2^n p$.

So, it suffices to prove the inequalities

$$\overline{\theta_1 \cdots \theta_{2^{n+1}p-i}} < \sigma^i(\theta_1 \cdots \theta^-_{2^n p}\theta_1 \cdots \theta_{2^n p}) < \theta_1 \cdots \theta_{2^{n+1}p-i} \tag{3.3}$$

for any $0 \le i < 2^n p$. By Lemma 2.5 it follows that for any $0 \le i < 2^n p$ we have

$$\overline{\theta_1 \cdots \theta_{2^n p-i}} \le \theta_{i+1} \cdots \theta^-_{2^n p} < \theta_1 \cdots \theta_{2^n p-i}$$

and

$$\theta_1 \cdots \theta_i > \overline{\theta_{2^n p-i+1} \cdots \theta_{2^n p}}.$$

This proves (3.3). □

Lemma 3.3. *Let* $(q_0, q_0^*) \subset (q', M+1) \setminus \overline{\mathcal{U}}$ *be a connected component generated by* $t_1 \cdots t_p$. *Then* $\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_{q_0^*}) > 0$ *for any* $r \in (q_0^*, M+1]$.

Proof. Take $r \in (q_0^*, M+1]$. By Lemma 3.1 there exists $n \ge 1$ such that

$$r_n \in (q_0^*, r) \cap \mathcal{U}.$$

Write $(\theta_i) = \alpha(q_0^*)$ and let $\xi_n = \theta_1 \cdots \theta_{2^n p}$. Denote by $X_A^{(n)}$ the subshift of finite type over the states $\{\xi_n, \xi_n^-, \overline{\xi_n}, \overline{\xi_n^-}\}$ with adjacency matrix

$$A = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Note that $\alpha(r_n) = \theta_1 \cdots \theta_{2^n p}(\theta_{2^n p+1} \cdots \theta_{2^{n+1}p})^\infty$. Then by Lemmas 3.2 and 2.3 it follows that

$$X_A^{(n)} \subseteq \mathcal{U}'_{r_n} \subseteq \mathcal{U}'_r. \tag{3.4}$$

Furthermore, note that

$$\begin{aligned}\xi_n\overline{\xi_n^-}(\overline{\xi_n}\xi_n)^3 &= \theta_1\cdots\theta_{2^{n+1}p}(\overline{\theta_1\cdots\theta_{2^{n+1}p}}^+)^3\\ &= \theta_1\cdots\theta_{2^{n+2}p}(\overline{\theta_1\cdots\theta_{2^{n+1}p}}^+)^2\\ &> \theta_1\cdots\theta_{2^{n+2}p}\overline{\theta_1\cdots\theta_{2^{n+1}p}\theta_{2^{n+1}p+1}\cdots\theta_{2^{n+2}p}}^+\\ &= \theta_1\cdots\theta_{2^{n+2}p}\theta_{2^{n+2}p+1}\cdots\theta_{2^{n+3}p}.\end{aligned}$$

Then by Lemmas 2.3 and 3.1 it follows that any sequence starting at

$$\mathbf{c} := \xi_n^-\xi_n\overline{\xi_n^-}(\overline{\xi_n}\xi_n)^3$$

can not belong to $\mathcal{U}'_{r_{n+2}}$. Therefore, by (3.4) we obtain

$$X_A^{(n)}(\mathbf{c}) := \left\{(d_i)\in X_A^{(n)} : d_1\cdots d_{(2^{n+3}+2^n)p}=\mathbf{c}\right\}\subseteq X_A^{(n)}\setminus\mathcal{U}'_{r_{n+2}}\subset\mathcal{U}'_r\setminus\mathcal{U}'_{q_0^*}. \tag{3.5}$$

Note that the subshift of finite type $X_A^{(n)}$ is irreducible (cf. [23]), and the image $\pi_{M+1}(X_A^{(n)})$ is a graph-directed set satisfying the open set condition (cf. [24]). Then by (3.5) it follows that

$$\begin{aligned}\dim_H\pi_{M+1}(\mathcal{U}'_r\setminus\mathcal{U}'_{q_0^*}) &\geq \dim_H\pi_{M+1}(X_A^{(n)}(\mathbf{c}))\\ &= \dim_H\pi_{M+1}(X_A^{(n)}) = \frac{\log\left((1+\sqrt{5})/2\right)}{2^n p\log(M+1)} > 0.\end{aligned}$$

□

The following lemma can be shown in a way which resembles closely the analysis in [22, Page 2829–2830]. For the sake of completeness we include a sketch of its proof.

Lemma 3.4. *Let $(q_0,q_0^*)\subset(q',M+1)\setminus\overline{\mathcal{U}}$ be a connected component. Then $\dim_H\pi_{M+1}(\mathcal{U}'_{q_0^*}\setminus\mathcal{U}'_{q_0})=0$.*

Sketch of the proof. Suppose that (q_0,q_0^*) is a connected component generated by $\mathbf{t}=t_1\cdots t_p$. Then

$$\alpha(q_0)=\mathbf{t}^\infty \quad\text{and}\quad \alpha(q_0^*)=\lim_{n\to\infty}g^n(\mathbf{t})=\mathbf{t}^+\,\overline{\mathbf{t}}\,\overline{\mathbf{t}^+}\,\mathbf{t}^+\cdots, \tag{3.6}$$

where $g(\cdot)$ is defined in (2.1).

For $n\geq 0$ let $\omega_n := g^n(\mathbf{t})^+$. Take $(d_i)\in\mathcal{U}'_{q_0^*}\setminus\mathcal{U}'_{q_0}$. Then by using (3.6) and Lemma 2.3 it follows that there exists $m\geq 1$ such that

$$\mathbf{t}^\infty=\alpha(q_0)\leq d_{m+1}d_{m+2}\cdots<\alpha(q_0^*)=\mathbf{t}^+\overline{\mathbf{t}}\cdots, \tag{3.7}$$

or symmetrically,

$$\mathbf{t}^\infty = \alpha(q_0) \le \overline{d_{m+1}d_{m+2}\cdots} < \alpha(q_0^*) = \mathbf{t}^+\overline{\mathbf{t}}\cdots. \tag{3.8}$$

Suppose $(d_{m+i}) \neq \mathbf{t}^\infty$ and $(d_{m+i}) \neq \overline{\mathbf{t}^\infty}$. Then there exists $u \ge m$ such that

$$d_{u+1}\cdots d_{u+p} = \mathbf{t}^+ = \omega_0 \quad \text{or} \quad d_{u+1}\cdots d_{u+p} = \overline{\mathbf{t}^+} = \overline{\omega_0}.$$

- If $d_{u+1}\cdots d_{u+p} = \omega_0 = \mathbf{t}^+$, then by (3.7) and Lemma 2.3 it follows that

$$d_{u+p+1}\cdots d_{u+2p} = \overline{\mathbf{t}^+} \quad \text{or} \quad d_{u+p+1}\cdots d_{u+2p} = \overline{\mathbf{t}}.$$

  This implies $d_{u+1}\cdots d_{u+2p} = \mathbf{t}^+\overline{\mathbf{t}^+} = \omega_0\,\overline{\omega_0}$ or $d_{u+1}\cdots d_{u+2p} = \mathbf{t}^+\overline{\mathbf{t}} = \omega_1$.
- If $d_{u+1}\cdots d_{u+p} = \overline{\omega_0} = \overline{\mathbf{t}^+}$, then by (3.8) and Lemma 2.3 it follows that

$$d_{u+p+1}\cdots d_{u+2p} = \mathbf{t}^+ \quad \text{or} \quad d_{u+p+1}\cdots d_{u+2p} = \mathbf{t}.$$

  This yields that $d_{u+1}\cdots d_{u+2p} = \overline{\omega_0}\,\omega_0$ or $d_{u+1}\cdots d_{u+2p} = \overline{\omega_1}$.

Note that for each $n \ge 0$ the word $g^n(\mathbf{t})^+\,\overline{g^n(\mathbf{t})}$ is a prefix of $\alpha(q_0^*)$. By iteration of the above arguments, one can show that if $d_{v+1}\cdots d_{v+2^np} = \omega_n$, then $d_{v+1}\cdots d_{v+2^{n+1}p} = \omega_n\overline{\omega_n}$ or ω_{n+1}. Symmetrically, if $d_{v+1}\cdots d_{v+2^np} = \overline{\omega_n}$, then $d_{v+1}\cdots d_{v+2^{n+1}p} = \overline{\omega_n}\omega_n$ or $\overline{\omega_{n+1}}$.

Hence, we conclude that (d_i) must end with

$$\mathbf{t}^*(\omega_{i_0}\overline{\omega_{i_0}})^*(\omega_{i_0}\overline{\omega_{j_0}})^{s_0}(\omega_{i_1}\overline{\omega_{i_1}})^*(\omega_{i_1}\overline{\omega_{j_1}})^{s_1}\cdots(\omega_{i_n}\overline{\omega_{i_n}})^*(\omega_{i_n}\overline{\omega_{j_n}})^{s_n}\cdots$$

or its reflections, where $s_n \in \{0,1\}$ and

$$0 = i_0 < j_0 \le i_1 < j_1 \le i_2 < \cdots \le i_n < j_n \le i_{n+1} < \cdots.$$

Here $*$ is an element of the set $\{0,1,2,\ldots\} \cup \{\infty\}$.

Since the length of $\omega_n = g^n(\mathbf{t})^+$ grows exponentially fast as $n \to \infty$, we conclude that $\dim_H \pi_{M+1}(\mathcal{U}'_{q_0^*} \setminus \mathcal{U}'_{q_0}) = 0$. □

Proof of Theorem 1.1 for (i) ⇔ (ii). First we prove (i) ⇒ (ii). If $q = q_0^*$ is the right endpoint of a connected component of $(q', M+1) \setminus \overline{\mathcal{U}}$, then by Lemma 3.3 we have

$$\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) > 0 \quad \text{for any} \quad r \in (q, M+1].$$

Clearly, it is trivial when $q = M+1$. Now we take $q \in (\mathcal{U} \setminus \{M+1\}) \setminus \bigcup\{q_0^*\}$ and take $r \in (q, M+1]$. By Lemma 2.6 (i) one can find $q_0^* \in (q, r)$, and therefore by Lemma 3.3 we obtain

$$\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) \ge \dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_{q_0^*}) > 0.$$

Now we prove (ii) $\Rightarrow$ (i). Take $q \in (1, M+1]\setminus\mathcal{U}$. We will show that $\dim_H \pi_{M+1}(\mathcal{U}'_r\setminus\mathcal{U}'_q) = 0$ for some $r \in (q, M+1]$. Note that $\bigcup\{q_0\} = \overline{\mathcal{U}}\setminus\mathcal{U}$. Then by (1.1) it follows that

$$q \in (1, q') \cup \bigcup [q_0, q_0^*).$$

Therefore, it suffices to prove $\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) = 0$ for some $r \in (q, M+1]$. We distinct the following two cases.

Case (I). $q \in (1, q')$. Then for any $r \in (q, q')$ we have

$$\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) \le \dim_H \pi_{M+1}(\mathcal{U}'_r) = 0,$$

where the last equality follows by [22, Theorem 4.6] (see also, [5, 15]).

Case (II). $q \in [q_0, q_0^*)$. Then for any $r \in (q, q_0^*)$ we have by Lemma 3.4 that

$$\dim_H \pi_{M+1}(\mathcal{U}'_r \setminus \mathcal{U}'_q) \le \dim_H \pi_{M+1}(\mathcal{U}'_{q_0^*} \setminus \mathcal{U}'_{q_0}) = 0.$$

□

3.2. **Proof of Theorem 1.1 for (i) $\Leftrightarrow$ (iii).** The following property for the Hausdorff dimension is well-known (cf. [14, Proposition 2.3]).

Lemma 3.5. *Let $f : (X, d_1) \to (Y, d_2)$ be a map between two metric spaces . If there exist constants $C > 0$ and $\lambda > 0$ such that*

$$d_2(f(x), f(y)) \le C d_1(x, y)^\lambda$$

for any $x, y \in X$, then $\dim_H X \ge \lambda \dim_H f(X)$.

Lemma 3.6. *Let $q \in \mathcal{U} \setminus \{M+1\}$. Then for any $r \in (q, M+1)$ we have*

$$\dim_H \mathcal{U} \cap (q, r) \ge \dim_H \pi_{M+1}\left(\{\alpha(p) : p \in \mathcal{U} \cap (q, r)\}\right).$$

Proof. Fix $q \in \mathcal{U} \setminus \{M+1\}$ and $r \in (q, M+1)$. Then Lemma 2.6 yields that $\mathcal{U} \cap (q, r)$ contains infinitely many elements. Take $p_1, p_2 \in \mathcal{U} \cap (q, r)$ with $p_1 < p_2$. Then by Lemma 2.1 we have $\alpha(p_1) < \alpha(p_2)$. So, there exists $n \ge 1$ such that

$$\alpha_1(p_1)\cdots\alpha_{n-1}(p_1) = \alpha_1(p_2)\cdots\alpha_{n-1}(p_2) \quad \text{and} \quad \alpha_n(p_1) < \alpha_n(p_2). \tag{3.9}$$

This implies

$$\begin{aligned} \pi_{M+1}(\alpha(p_2)) - \pi_{M+1}(\alpha(p_1)) &= \sum_{i=1}^{\infty} \frac{\alpha_i(p_2) - \alpha_i(p_1)}{(M+1)^i} \\ &\le \sum_{i=n}^{\infty} \frac{M}{(M+1)^i} = (M+1)^{1-n}. \end{aligned} \tag{3.10}$$

Note that $r < M+1$. By Lemma 2.1 we have $\alpha(r) < \alpha(M+1) = M^\infty$. Then there exists $N \geq 1$ such that

$$\alpha_1(r)\cdots\alpha_N(r) < \underbrace{M\cdots M}_{N}.$$

Therefore, by (3.9) and Lemma 2.3 we obtain

$$\sum_{i=1}^{n}\frac{\alpha_i(p_2)}{p_1^i} \geq \sum_{i=1}^{\infty}\frac{\alpha_i(p_1)}{p_1^i} = 1 = \sum_{i=1}^{\infty}\frac{\alpha_i(p_2)}{p_2^i} > \sum_{i=1}^{n}\frac{\alpha_i(p_2)}{p_2^i} + \frac{1}{p_2^{n+N}}.$$

Note that p_1, p_2 are elements of $\mathcal{U}$. Then $p_2 > p_1 \geq q'$. This implies

$$\begin{aligned}\frac{1}{(M+1)^{n+N}} < \frac{1}{p_2^{n+N}} &< \sum_{i=1}^{n}\left(\frac{\alpha_i(p_2)}{p_1^i} - \frac{\alpha_i(p_2)}{p_2^i}\right)\\ &\leq \sum_{i=1}^{\infty}\left(\frac{M}{p_1^i} - \frac{M}{p_2^i}\right) = \frac{M(p_2-p_1)}{(p_1-1)(p_2-1)} \leq \frac{M(p_2-p_1)}{(q'-1)^2}.\end{aligned}$$

Therefore, by (3.10) it follows that

$$\pi_{M+1}(\alpha(p_2)) - \pi_{M+1}(\alpha(p_1)) \leq (M+1)^{1-n} \leq \frac{(M+1)^{2+N}}{(q'-1)^2}(p_2-p_1).$$

Furthermore, by Lemma 2.1 it follows that $\pi_{M+1}(\alpha(p_2)) - \pi_{M+1}(\alpha(p_1)) \geq 0$. Hence, by using

$$f = \pi_{M+1}\circ\alpha: \quad \mathcal{U}\cap(q,r) \to \pi_{M+1}(\{\alpha(p) : p \in \mathcal{U}\cap(q,r)\})$$

in Lemma 3.5 we establish the lemma. □

Lemma 3.7. *Let (q_0, q_0^*) be a connected component of $(q', M+1)\setminus\overline{\mathcal{U}}$. Then $\dim_H \mathcal{U}\cap(q_0^*, r) > 0$ for any $r \in (q_0^*, M+1]$.*

Proof. Suppose that (q_0, q_0^*) is a connected component generated by $t_1\cdots t_p$. Let $(\theta_i) = \alpha(q_0^*)$. For $n \geq 2$ we write $\xi_n = \theta_1\cdots\theta_{2^n p}$, and denote by

$$\Gamma_n' := \left\{(d_i) : d_1\cdots d_{2^{n+1}p} = \xi_{n-1}(\overline{\xi_{n-1}}^+)^3, \quad (d_{2^{n+1}p+i}) \in X_A^{(n)}(\overline{\xi_n})\right\}.$$

Here $X_A^{(n)}(\overline{\xi_n})$ is the follower set of $\overline{\xi_n}$ in the subshift of finite type $X_A^{(n)}$ defined in (3.5). Now we claim that any sequence $(d_i) \in \Gamma_n'$ satisfies

$$\overline{(d_i)} < \sigma^j((d_i)) < (d_i) \quad \text{for all} \quad j \geq 1. \tag{3.11}$$

Take $(d_i) \in \Gamma_n'$. Then we deduce by the definition of Γ_n' that

$$d_1\cdots d_{2^{n+1}p+2^{n-1}p} = \theta_1\cdots\theta_{2^{n-1}p}(\overline{\theta_1\cdots\theta_{2^{n-1}p}}^+)^3\,\overline{\theta_1\cdots\theta_{2^n p}}. \tag{3.12}$$

We will split the proof of (3.11) into the following five cases.

(a) $1 \le j < 2^{n-1}p$. By (3.12) and Lemma 2.5 it follows that

$$\overline{\theta_1 \cdots \theta_{2^{n-1}p-j}} < d_{j+1} \cdots d_{2^{n-1}p} = \theta_{j+1} \cdots \theta_{2^{n-1}p} \le \theta_1 \cdots \theta_{2^{n-1}p-j},$$

and

$$d_{2^{n-1}p+1} \cdots d_{2^{n-1}p+j} = \overline{\theta_1 \cdots \theta_j} < \theta_{2^{n-1}p-j+1} \cdots \theta_{2^{n-1}p}.$$

This implies that (3.11) holds for all $1 \le j < 2^{n-1}p$.

(b) $2^{n-1}p \le j < 2^n p$. Let $k = j - 2^{n-1}p$. Then $0 \le k < 2^{n-1}p$. Clearly, if $k = 0$, then by using $\theta_1 > \overline{\theta_1}$ and $n \ge 2$ it yields that

$$\overline{\theta_1 \cdots \theta_{2^{n-1}p}} < d_{j+1} \cdots d_{2^n p} = \overline{\theta_1 \cdots \theta_{2^{n-1}p}}^{+} < \theta_1 \cdots \theta_{2^{n-1}p}.$$

Now we assume $1 \le k < 2^{n-1}p$. Then by (3.12) and Lemma 2.5 it follows that

$$\overline{\theta_1 \cdots \theta_{2^{n-1}p-k}} < d_{j+1} \cdots d_{2^n p} = \overline{\theta_{k+1} \cdots \theta_{2^{n-1}p}}^{+} \le \theta_1 \cdots \theta_{2^{n-1}p-k},$$

and

$$d_{2^n p+1} \cdots d_{2^n p+k} = \overline{\theta_1 \cdots \theta_k} < \theta_{2^{n-1}p-k+1} \cdots \theta_{2^{n-1}p}.$$

Therefore, (3.11) holds for all $2^{n-1}p \le j < 2^n p$.

(c) $2^n p \le j < 2^n p + 2^{n-1}p$. Let $k = j - 2^n p$. Then in a similar way as in Case (b) one can prove (3.11).

(d) $2^n p + 2^{n-1}p \le j < 2^{n+1}p$. Let $k = j - 2^n p - 2^{n-1}p$. Again by the same arguments as in Case (b) we obtain (3.11).

(e) $j \ge 2^{n+1}p$. Note that

$$d_1 \cdots d_{2^{n+1}p} = \theta_1 \cdots \theta_{2^{n-1}p}(\overline{\theta_1 \cdots \theta_{2^{n-1}p}}^{+})^3 > \theta_1 \cdots \theta_{2^{n+1}p}.$$

Then (3.11) follows by Lemma 3.2.

Therefore, by (3.11) and Lemma 2.4 it follows that any sequence in Γ'_n corresponds to a unique base $q \in \mathcal{U}$. Furthermore, by (3.12) and Lemma 3.1 each sequence $(d_i) \in \Gamma'_n$ satisfies

$$\alpha(q_0^*) = (\theta_i) < (d_i) < \theta_1 \cdots \theta_{2^{n-1}p}(\overline{\theta_1 \cdots \theta_{2^{n-1}p}}^{+})^{\infty} = \alpha(r_{n-1}).$$

Then by Lemma 2.1 it follows that

$$\alpha(q) \in \Gamma'_n \quad \Longrightarrow \quad q \in \mathcal{U} \cap (q_0^*, r_{n-1}).$$

Fix $r > q_0^*$. So by Lemma 3.1 there exists a sufficiently large integer $n \ge 2$ such that

$$\Gamma'_n \subset \{\alpha(q) : q \in \mathcal{U} \cap (q_0^*, r)\}. \tag{3.13}$$

Note by the proof of Lemma 3.3 that $X_A^{(n)}$ is an irreducible subshift of finite type over the states $\{\xi_n, \xi_n^-, \overline{\xi_n}, \overline{\xi_n^-}\}$. Hence, by (3.13) and Lemma 3.6 it follows that

$$\begin{aligned}\dim_H \mathcal{U} \cap (q_0^*, r) \geq \dim_H \pi_{M+1}(\Gamma_n') &= \dim_H \pi_{M+1}(X_A^{(n)}) \\ &= \frac{\log\left((1+\sqrt{5})/2\right)}{2^n p \log(M+1)} > 0.\end{aligned}$$

□

Proof of Theorem 1.1 for (i) ⇔ (iii). First we prove (i) ⇒ (iii). Excluding the trivial case $q = M+1$ we take $q \in \mathcal{U} \setminus \{M+1\}$. Suppose that $r \in (q, M+1]$. If $q = q_0^*$, then by Lemma 3.7 we have $\dim_H \mathcal{U} \cap (q, r) > 0$.

If $q \in (\mathcal{U} \setminus \{M+1\}) \setminus \bigcup\{q_0^*\}$, then by Lemma 2.6 (i) there exists $q_0^* \in (q, r)$. So, by Lemma 3.7 we have

$$\dim_H \mathcal{U} \cap (q, r) \geq \dim_H \mathcal{U} \cap (q_0^*, r) > 0.$$

Now we prove (iii) ⇒ (i). Suppose on the contrary that $q \in (1, M+1] \setminus \mathcal{U}$. We will show that $\mathcal{U} \cap (q, r) = \emptyset$ for some $r \in (q, M+1]$. Take $q \in (1, M+1] \setminus \mathcal{U}$. By (1.1) it follows that

$$q \in (1, q') \cap \bigcup [q_0, q_0^*).$$

This implies that $\mathcal{U} \cap (q, r) = \emptyset$ for $r \in (q, M+1]$ sufficiently close to q. □

3.3. **Proof of Theorem 1.2.**

Proof of Theorem 1.2. (i) ⇒ (ii). Take $q \in \overline{\mathcal{U}} \setminus (\bigcup\{q_0^*\} \cup \{q'\})$ and $p \in (1, q)$. By Lemma 2.6 (ii) there exists $q_0^* \in (p, q)$. Hence, by Lemma 3.3 it follows that

$$\dim_H \pi_{M+1}(\mathcal{U}_q' \setminus \mathcal{U}_p') \geq \dim_H \pi_{M+1}(\mathcal{U}_q' \setminus \mathcal{U}_{q_0^*}') > 0.$$

(ii) ⇒ (i). Suppose on the contrary that $q \notin \overline{\mathcal{U}} \setminus (\bigcup\{q_0^*\} \cup \{q'\})$. Then by (1.1) we have

$$q \in (1, q'] \cup \bigcup (q_0, q_0^*].$$

By using Lemma 3.4 it follows that for $p \in (1, q)$ sufficiently close to q we have $\dim_H \pi_{M+1}(\mathcal{U}_q' \setminus \mathcal{U}_p') = 0$.

(i) ⇒ (iii). Take $q \in \overline{\mathcal{U}} \setminus (\bigcup\{q_0^*\} \cup \{q'\})$ and $p \in (1, q)$. By Lemma 2.6 (ii) there exists $q_0^* \in (p, q)$. Hence, by Lemma 3.7 it follows that

$$\dim_H \mathcal{U} \cap (p, q) \geq \dim_H \mathcal{U} \cap (q_0^*, q) > 0.$$

(iii) ⇒ (i). Suppose $q \notin \overline{\mathcal{U}} \setminus (\bigcup\{q_0^*\} \cup \{q'\})$. Then by (1.1) we have $q \in (1, q'] \cup \bigcup (q_0, q_0^*]$. So, for $p \in (1, q)$ sufficiently close to q we have $\mathcal{U} \cap (p, q) = \emptyset$. □

Acknowledgements

The authors thank the anonymous referees for many useful comments. The first author was supported by NSFC No. 11401516 and Jiangsu Province Natural Science Foundation for the Youth No. BK20130433. The second author was supported by NSFC No. 11271137, 11571144, 11671147 and Science and Technology Commission of Shanghai Municipality (STCSM) No. 13dz2260400. The third author was supported by NSFC No. 11601358.

References

[1] Allouche J.-P., Cosnard, M.: Itérations de fonctions unimodales et suites engendrées par automates. C. R. Acad. Sci. Paris Sér. I Math. **296**, 159–162 (1983)

[2] Allouche, J.-P., Frougny, C., Hare, K. G.: On univoque Pisot numbers. Math. Comp. **76**, 1639–1660 (2007)

[3] Allouche J.-P., Shallit, J.: The ubiquitous Prouhet-Thue-Morse sequence, Sequences and their applications (Singapore, 1998), Springer Ser. Discrete Math. Theor. Comput. Sci., Springer, London, pp. 1–16 (1999)

[4] Baiocchi C., Komornik, V.: Greedy and quasi-greedy expansions in non-integer bases, arXiv:0710.3001v1 (2007).

[5] Baker, S.: Generalized golden ratios over integer alphabets. Integers **14**, 28 pages (2014)

[6] Daróczy Z., Kátai, I.: Univoque sequences. Publ. Math. Debrecen **42**, 397–407 (1993)

[7] Daróczy Z., Kátai, I.: On the structure of univoque numbers. Publ. Math. Debrecen **46**, 385–408 (1995)

[8] de Vries, M.: On the number of unique expansions in non-integer bases. Topology and its Applications **156**, 652–657 (2009)

[9] de Vries, M, Komornik, V.: Unique expansions of real numbers, Adv. Math. **221**, 390–427 (2009)

[10] de Vries, M., Komornik, V., Loreti, P.: Topology of the set of univoque bases. Topology and its Applications **205**, 117–137 (2016)

[11] Erdős, P., Joó, I., Komornik, V.: Characterization of the unique expansions $1 = \sum_{i=1}^{\infty} q^{-n_i}$ and related problems. Bull. Soc. Math. France **118**, 377–390 (1990)

[12] Erdős, P., Horváth, M., Joó, I.: On the uniqueness of the expansions $1 = \sum q^{-n_i}$. Acta Math. Hungar. **58**, 333–342 (1991)

[13] Erdős P., Joó, I.: On the number of expansions $1 = \sum q^{-n_i}$. Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **35**, 129–132 (1992)

[14] Falconer, K.: Fractal geometry. John Wiley & Sons Ltd., Chichester, Mathematical foundations and applications (1990)

[15] Glendinning, P., Sidorov, N.: Unique representations of real numbers in non-integer bases. Math. Res. Lett. **8**, 535–543 (2001)

[16] Komornik, V.: Expansions in noninteger bases. Integers **11B**, 30 pages (2011)

[17] Komornik, V., Kong, D., Li, W.: Hausdorff dimension of univoque sets and Devil's staircase. Adv. Math. **305**, 165–196 (2017)

[18] Komornik, V., Loreti, P.: Unique developments in non-integer bases. Amer. Math. Monthly **105**, 636–639 (1998)
[19] Komornik, V., Loreti, P.: Subexpansions, superexpansions and uniqueness properties in non-integer bases. Period. Math. Hungar. **44**, 197–218 (2002)
[20] Komornik, V., Loreti, P.: On the topological structure of univoque sets. J. Number Theory **122**, 157–183 (2007)
[21] Kong, D., Li, W.: Hausdorff dimension of unique beta expansions. Nonlinearity **28**, 187–209 (2015)
[22] Kong, D., Li, W., Dekking, M.: Intersections of homogeneous Cantor sets and beta-expansions. Nonlinearity **23**, 2815–2834 (2010)
[23] Lind, D., Marcus, B.: An introduction to symbolic dynamics and coding. Cambridge University Press, Cambridge (1995)
[24] Mauldin, R.D., Williams, S.C.: Hausdorff dimension in graph directed constructions. Trans. Amer. Math. Soc. **309**, 811–829 (1988)
[25] Parry, W.: On the β-expansions of real numbers. Acta Math. Acad. Sci. Hungar. **11**, 401–416 (1960)
[26] Rényi, A.: *Representations for real numbers and their ergodic properties*. Acta Math. Acad. Sci. Hungar. **8**, 477–493 (1957)
[27] Sidorov, N.: Expansions in non-integer bases: lower, middle and top orders. J. Number Theory **129**, 741–754 (2009)

School of Mathematical Science, Yangzhou University, Yangzhou, JiangSu 225002, People's Republic of China

Current address: Mathematical Institute, University of Leiden, PO Box 9512, 2300 RA Leiden, The Netherlands

E-mail address: derongkong@126.com

Department of Mathematics, Shanghai Key Laboratory of PMMP, East China Normal University, Shanghai 200062, People's Republic of China

E-mail address: wxli@math.ecnu.edu.cn

Department of Mathematics, Sichuan Normal University, Chengdu 610068, People's Republic of China

E-mail address: lvfan1123@163.com

Tussen de Grachten 213, 1381 DZ, Weesp, The Netherlands

E-mail address: martijndevries0@gmail.com